

## Investigation 1: How Loud is too Loud?

## Decibels and Sound Intensity

**Think About This Situation**

Study the sound intensity values, sources, and decibel ratings given in the table.

a) Are the intensity and decibel numbers in the order of loudness that you would expect for the different familiar sources?

b) What pattern do you see relating the sound intensity values (watts/cm<sup>2</sup>) and the decibel numbers?

Use the table below to answer the Think About This Situation questions.

Answer a) \_\_\_\_\_

Answer b) \_\_\_\_\_

Sound Intensity (in watts/cm <sup>2</sup> )	Noise Source	Relative Intensity (in decibels)
$10^3$	Military rifle	150
$10^2$	Jet plane (30 meters away)	140
$10^1$	[Level at which sound is painful]	130
$10^0$	Amplified rock music	120
$10^{-1}$	Power tools	110
$10^{-2}$	Noisy kitchen	100
$10^{-3}$	Heavy traffic	90
$10^{-4}$	Traffic noise in a small car	80
$10^{-5}$	Vacuum cleaner	70
$10^{-6}$	Normal conversation	60
$10^{-7}$	Average home	50
$10^{-8}$	Quiet conversation	40
$10^{-9}$	Soft whisper	30
$10^{-10}$	Quiet living room	20
$10^{-11}$	Quiet recording studio	10
$10^{-12}$	[Barely audible]	0

Source: *Real-Life Math: Everyday Use of Mathematical Concepts*, Evan Glazer and John McConnell, 2002.

Your analysis of the sound intensity data might have suggested several different algorithms for converting watts per cm<sup>2</sup> into decibels. For example, **if the intensity of a sound is  $10^x$  watts/cm<sup>2</sup>, its loudness in decibels is  $10x + 120$ .**

The key to discovery of this conversion rule is the fact that all sound intensities were written as powers of 10. What would you have done if the sound intensity readings had been written as number like 3.45 watts/cm<sup>2</sup> or 0.0023 watts/cm<sup>2</sup>?

As you work on problems of this investigation, look for an answer to this question:

How can any positive number be expressed as a power of 10?

- 1) Express each of the numbers below as accurately as possible as a power of 10. You can find exact values for some of the required exponents by thinking about the meanings of positive and negative exponents. Others might require some calculator exploration of ordered pairs that satisfy the exponential equation  $y = 10^x$ .
- |          |           |              |
|----------|-----------|--------------|
| a. 100   | b. 10,000 | c. 1,000,000 |
| d. 0.01  | e. -0.001 | f. 3.45      |
| g. -34.5 | h. 345    | i. 0.0023    |
- 2) Suppose that the sound intensity of a screaming baby was measured as 9.5 watts/cm<sup>2</sup>. To calculate the equivalent intensity in decibels, 9.5 must be written as 10<sup>x</sup> for some value of x.
- Between which two integers does it make sense to look for values of the required exponent? How do you know?
  - Which of the two integer values in Part a is probably closer to the required power of 10?
  - Estimate the required exponent to the nearest hundredth. Then use your estimate to calculate a decibel rating for the loudness of the baby's scream.
  - Estimate the decibel rating for loudness of sound from a television set that registers intensity of 6.2 watts/cm<sup>2</sup>.

The exponents in the example above demonstrate logarithms. A **logarithm** is defined as follows:

The logarithm base  $b$  of a positive number  $y$  is defined as follows:

$$\text{If } y = b^x, \text{ then } \log_b y = x.$$

The exponent  $x$  in the exponential expression  $b^x$  is the logarithm in the equation  $\log_b y = x$ . The base  $b$  in  $b^x$  is the same as the base  $b$  in the logarithm. In both cases,  $b \neq 1$  and  $b > 0$ . So what this means is that you use logarithms to undo exponential expressions or equations and you use exponents to undo logarithms, which means that the operations are inverses of each other. Thus, an exponential function is the inverse of a logarithmic function and vice versa.

### Common Logarithms

As you discovered in your work on Problems 1 and 2, it is not easy to solve equations like  $10^x = 9.5$  or  $10^x = 0.0023$ , even by estimation. To deal with this very important problem, mathematicians have developed procedures for finding exponents. We have already discussed logarithms in the previous lesson, but in this lesson we will solve exponent problems which focus on a base of 10. When we solve logarithmic problems in base 10, we call them **common logarithms**.

The definition of common logarithms is usually expressed as:

$$\log_{10} a = b \text{ if and only if } 10^b = a$$

$\log_{10} a$  is pronounced “log base 10 of  $a$ ”. Because base 10 logarithms are so commonly used,  $\log_{10} a$  is often written as  $\log a$ . Most calculators have a built-in log function that automatically finds the required exponent value.

- 3) Use your calculator to find the following logarithms. Then compare the results with your work on Problem 1.
- |                   |                    |                     |
|-------------------|--------------------|---------------------|
| a. $\log 100$     | b. $\log 10,000$   | c. $\log 1,000,000$ |
| d. $\log 0.01$    | e. $\log (-0.001)$ | f. $\log 3.45$      |
| g. $\log (-3.45)$ | h. $\log 345$      | i. $\log 0.0023$    |
- 4) What do your results from Problem 3 (especially Parts e and g) suggest about the kinds of numbers that have logarithms? See if you can explain your answer by using the connection between logarithms and the exponential function  $y = 10^x$ .
- 5) Logarithms can be used to calculate the decibel rating of sounds, when the intensity is measured in  $\text{watts/cm}^2$ .
- a. Use the logarithm feature of your calculator to rewrite 9.5 as a power of 10. That is, find  $x$  so that  $9.5 = 10^x$ .
- b. Recall that if the intensity of a sound is  $10^x \text{ watts/cm}^2$ , then the expression  $10x + 120$  can be used to convert the sound’s intensity to decibels. Use your results from Part a to find the decibel rating of the crying baby in Problem 2.
- 6) Assume the intensity of a sound  $I = 10^x \text{ watts/cm}^2$ .
- a. Explain why  $x = \log I$ .
- b. Rewrite the expression for converting sound intensity reading to decibel numbers using  $\log I$ .
- 7) Use your conversion expression from Problem 6 to find the decibel rating of the television set in Problem 2 Part d.

### Why Do They Taste Different?

You may recall from your study of science that the acidity of a substance is described by the pH rating – the lower the pH rating, the more acidic the substance is. The acidity depends upon the hydrogen ion concentration in the substance (in moles per liter). Some sample hydrogen ion concentrations are given below. Since those hydrogen ion concentrations are generally very small numbers, they are converted to the simpler pH scale for reporting.

8) Examine the table at right.

a. Describe how hydrogen ion concentrations  $[H^+]$  are into pH readings.

Substance	$[H^+]$	pH
Hand soap	$10^{-10}$	10
Egg white	$10^{-9}$	9
Sea water	$10^{-8}$	8
Pure water	$10^{-7}$	7
White bread	$10^{-6}$	6
Coffee	$10^{-5}$	5
Tomato juice	$10^{-4}$	4
Orange juice	$10^{-3}$	3

converted

b. Write an equation that makes use of logarithms expressing function of hydrogen ion concentration  $[H^+]$ .

pH as a

9) Use the equation relating hydrogen ion concentration and pH reading to compare acidity of some familiar liquids.

a. Complete the table at right. Round results to the

Substance	$[H^+]$ Proportion	pH Reading
lemonade	0.00501	
apple juice	0.000794	
milk	0.000000355	

nearest tenth.

b. Explain how your results tell which is more acidic – apple juice, or milk.

lemonade,

## Summarize the Mathematics

In work on the problems of this investigation, you learned how physical measurements of sound intensity and acidity of a chemical substance are converted into the more familiar decibel and pH numbers. You also learned how the *logarithm* function is used in those processes.

- a. How would you explain to someone who did not know about logarithms what the expression  $\log y = x$  tells about the numbers  $x$  and  $y$ ?
- b. What can be said about the value of  $\log y$  in each case below? Give brief justifications of your answers.
- $0 < y < 1$
  - $1 < y < 10$
  - $10 < y < 100$
  - $100 < y < 1,000$

*Be prepared to explain your ideas to the class.*

A: \_\_\_\_\_

Bi. \_\_\_\_\_

Bii. \_\_\_\_\_

Biii. \_\_\_\_\_

Biv. \_\_\_\_\_

### Check Your Understanding

Use your understanding of the relationship between logarithms and exponents to help complete these tasks.

- a. Find these common (base 10) logarithms without using a calculator.
- $\log 1,000$
  - $\log 0.001$
  - $\log 10^{3.2}$
- b. Use the function  $y = 10^x$ , but not the logarithm key of your calculator to estimate each of these logarithms to the nearest tenth. Explain how you arrived at your answers.
- $\log 75$
  - $\log 750$
  - $\log 7.5$
- c. If the intensity of sound from a drag race car is  $125 \text{ watts/cm}^2$ , what is the decibel rating of the loudness for that sound?